



Endogenous brassinosteroids in microalgae exposed to salt and low temperature stress

Wendy A. Stirk, Péter Bálint, Danuše Tarkowská, Miroslav Strnad, Johannes van Staden & Vince Ördög

To cite this article: Wendy A. Stirk, Péter Bálint, Danuše Tarkowská, Miroslav Strnad, Johannes van Staden & Vince Ördög (2018) Endogenous brassinosteroids in microalgae exposed to salt and low temperature stress, *European Journal of Phycology*, 53:3, 273-279, DOI: [10.1080/09670262.2018.1441447](https://doi.org/10.1080/09670262.2018.1441447)

To link to this article: <https://doi.org/10.1080/09670262.2018.1441447>



Published online: 24 Apr 2018.



Submit your article to this journal [↗](#)



Article views: 487



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 8 View citing articles [↗](#)

Endogenous brassinosteroids in microalgae exposed to salt and low temperature stress

Wendy A. Stirk^a, Péter Bálint^b, Danuše Tarkowská^c, Miroslav Strnad^c, Johannes van Staden^a and Vince Ördög^{a,b}

^aResearch Centre for Plant Growth and Development, School of Life Sciences, University of KwaZulu-Natal, Pietermaritzburg, P/Bag X 01, Scottsville 3209, South Africa; ^bDepartment of Plant Sciences, Faculty of Agricultural and Food Sciences, Széchenyi István University, Kolbai K. Str. 8, H-9200 Mosonmagyaróvár, Hungary; ^cLaboratory of Growth Regulators, Centre of the Region Haná for Biotechnological and Agricultural Research, Palacký University & Institute of Experimental Botany AS CR, Šlechtitelů 27, Olomouc CZ-783 71, Czech Republic

ABSTRACT

Brassinosteroids are part of the hormonal network that regulates growth processes and stress responses in plants. There is evidence for a similar hormonal network in microalgae. In the present study, six microalgae (*Chlorococcum ellipsoideum*, *Gyoeffya humicola*, *Nautococcus mamillatus*, *Acutodesmus acuminatus*, *Protococcus viridis* and *Chlorella vulgaris*) were subjected to salt and low temperature stress with the addition of 36 g l⁻¹ NaCl and transfer from 25°C to 15°C. There was a rapid response to salt stress with the brassinosteroid content (mainly castasterone with lower amounts of brassinolide, homocastasterone and typhasterol) increasing within 30 min of the salt treatment and remaining at these elevated levels after 7 h. The decrease in temperature had little effect on the brassinosteroid content. This was the first study to show that endogenous brassinosteroids increase in response to abiotic stress in a number of microalgae species.

ARTICLE HISTORY Received 5 October 2017; Revised 12 January 2018; Accepted 16 January 2018

KEY WORDS Brassinolide; castasterone; homocastasterone; microalgae; salt stress; typhasterol

Introduction

Brassinosteroids are polyhydroxylated steroid hormones with a ubiquitous distribution in the Plant Kingdom, occurring in angiosperms, gymnosperms (Bajguz & Tretyn, 2003) and non-seed plants, e.g. in the strobilus of *Equisetum arvense* (pteridophyte; Takatsuto *et al.*, 1990) and *Marchantia polymorpha* (bryophyte; Park *et al.*, 1999). They are also present in seaweed, e.g. *Ecklonia maxima* (Phaeophyceae; Stirk *et al.*, 2014), freshwater Chlorophyceae, e.g. *Hydrodictyon reticulatum* (Hydrodictyaceae; Yokota *et al.*, 1987) and microalgae (Bajguz, 2009a; Stirk *et al.*, 2013a).

Brassinosteroids consist of a 5 α -cholestane skeleton with structural variations in the type and position of the A- and B- rings and side chain (Oklestková *et al.*, 2015). Variations include the number of hydroxyl (OH) groups on the A-ring, the oxidation stage of the B-ring (7-oxalactone, 6-ketone and non B-ring oxidized types) and substituents at mainly C-22, C-23 and C-24 on the side chain (Bajguz & Tretyn, 2003). There are two interconnected biosynthetic pathways for C₂₈ brassinosteroids, namely the early C-6 oxidation and late C-6 oxidation pathways, with campersterol the precursor for both pathways (Fujioka & Yokota, 2003). C₂₉ brassinosteroids follow a different biosynthetic pathway beginning with the precursor sitosterol and C₂₇

brassinosteroids are synthesized from the precursor cholesterol (Fujioka & Yokota, 2003; Oklestková *et al.*, 2015). While over 70 brassinosteroid metabolites have been identified to date (Bajguz & Tretyn, 2003; Bajguz, 2009a), only a few are biologically active. Brassinolide (BL) and its direct biosynthetic precursor castasterone (CS) and 24-*epi*brassinolide show the highest biological activity (Bartwell *et al.*, 2013). Levels of active brassinosteroids are regulated by their modification mainly via hydroxylation, oxidation and epimerization reactions, and conjugation with either glucose and/or fatty acids (Fujioka & Yokota, 2003).

Brassinosteroids regulate many physiological and developmental processes in plants. They promote cell division, expansion and differentiation at the cellular level and influence root and shoot growth, reproductive processes and seed germination at the whole plant level (Fariduddin *et al.*, 2014; Oklestková *et al.*, 2015; Rajewska *et al.*, 2016). There is evidence that brassinosteroids have a similar regulatory role in microalgae. Exogenous application of two brassinosteroids i.e. BL and CS stimulated cell division, increased net photosynthetic rate and chlorophyll content and increased monosaccharide and organic and inorganic phosphorus content in *Chlorella vulgaris* (Bajguz & Czerpak, 1998). Brassinazole, an inhibitor of brassinosteroid biosynthesis, suppressed the growth of *C. vulgaris* with a decrease in RNA synthesis,

and protein, carotenoid and sugar content. This inhibitory effect was partially reversed with the co-application of BL (Bajguz & Asami, 2004).

Brassinosteroids also play an important role in alleviating abiotic stresses in plants through interactions with the hormonal network and by stimulating the antioxidant system, modifying both antioxidant enzymes and non-enzymatic antioxidants in plants (Fariduddin *et al.*, 2014; Oklestková *et al.*, 2015; Rajewska *et al.*, 2016). For example, brassinosteroids increased antioxidant enzyme activity and carotenoid content in drought-stressed maize seedlings (Fariduddin *et al.*, 2014); brassinosteroids applied to salinity-stressed rice, *Arabidopsis thaliana*, *Brassica napus*, *Vigna radiata* and *Cicer arietinum* modified the accumulation of proline and the activity of various antioxidant enzymes and altered gene expression (reviewed in Fariduddin *et al.*, 2014) and brassinosteroids improved the tolerance and recovery of tomato seedlings to high temperatures by increasing the accumulation of heat-shock proteins as well as providing protection from cold stress (Fariduddin *et al.*, 2014). Similar responses occurred in *Chlorella vulgaris* where exogenous application of BL partially overcame the inhibitory effects of heavy metal stress on growth, this was accompanied by an increase in both enzymatic (superoxide dismutase, catalase, ascorbate peroxidase and glutathione reductase) and non-enzymatic (ascorbate and glutathione) systems in a concentration-dependent manner (Bajguz, 2010).

There is interest in using cell culture techniques for the *in vivo* production of bioactive plant metabolites as it allows for a continuous and reliable source of natural products (Bartwell *et al.*, 2013). Brassinosteroids have promising therapeutic properties such as antiviral (herpes, measles, polio), antimicrobial, anti-angiogenic, anti-proliferative (anti-cancer) and neuroprotective (antioxidant) activities (reviewed in Bajguz *et al.*, 2013; Oklestková *et al.*, 2015) combined with a favourable safety profile with low toxicity (Bajguz *et al.*, 2013). Due to similarities between brassinosteroids and animal steroids, with both being cholesterol derivatives, brassinosteroids have potential to provide novel leads for drug development. Understanding the production of brassinosteroids in microalgae in response to induced stress can enhance the value of the biomass.

The aim of the present study was to quantify changes in the endogenous brassinosteroid content in microalgae exposed to salt and cold temperature stress.

Materials and methods

Microalgae cultures, growth conditions and experimental design

Six axenic microalgae strains from different taxonomic groups were selected from the Mosonmagyaróvár Algal Culture Collection (MACC) for the study. Details of the

Table 1. Taxonomic details and culture origin of the six microalgae strains used in the study.

Class	Genus and species	MACC	Origin
Chlorophyceae	<i>Chlorococcum ellipsoideum</i> Deason & Bold	712	Czech Republic, soil
	<i>Gyoefferfyaana humicola</i> Kol & Chodat	334	Brazil, soil
	<i>Nautococcus mamillatus</i> Korschikov	716	Czech Republic, soil
	<i>Acutodesmus acuminatus</i> ^a (Lagerh.) Tsarenko	41	Czech Republic, soil
	<i>Protococcus viridis</i> C. Agardh	219	Brazil, soil
Trebouxiophyceae	<i>Chlorella vulgaris</i> Beyerinck	755	Czech Republic, soil

^aFormerly *Scenedesmus acuminatus*.

strains are shown in Table 1. Each strain was inoculated from agar cultures into two flasks containing 250 ml liquid Bristol nutrient medium (Bold, 1949). The cultures were grown at $25 \pm 2^\circ\text{C}$ in a 14:10 h light:dark photoperiod and illuminated from below with $130 \mu\text{mol photons m}^{-2} \text{s}^{-1}$ and aerated with 20 l h^{-1} 1.5% CO_2 -enriched sterile air during the light phase.

After 7 days, the suspension cultures from both flasks were combined and used to inoculate 24 flasks each containing 250 ml Bristol medium. The starting biomass was 10 mg l^{-1} . These cultures were grown in the above-mentioned culture conditions for a further 4 days.

At the beginning of the light period (7.00 am) on day 4, the flasks were divided into four groups (6 flasks per treatment) and treated as follows:

- (i) 30 ml distilled water added and flasks incubated at 25°C ;
- (ii) 30 ml sterile NaCl solution added to give a concentration of 36 g l^{-1} NaCl and flasks incubated at 25°C ;
- (iii) 30 ml distilled water added and flasks incubated at 15°C ;
- (iv) 30 ml sterile NaCl solution added to give a concentration of 36 g l^{-1} NaCl and flasks incubated at 15°C .

After 30 min and again at 7 h (2.00 pm), 10 ml samples were collected from the flasks for each treatment. A portion from each sample (2 ml) was stored deep-frozen for brassinosteroid quantification and another 2 ml were preserved with Lugol's solution for cell number counting. The remaining sample (5 ml) was used for dry weight determination.

Cell number

Cell number was counted using an Olympus BX60 microscope (Olympus Optical Co. Ltd, Tokyo, Japan)

and analysed with analySYS software (Software Imaging GmbH, Münster, Germany). There were six replicates per treatment. One-way ANOVA followed by *post hoc* Tukey's test were used to detect significant differences ($P > 0.05$) between the treatments (SigmaPlot v.13).

Dry weight determination

Samples were filtered through Whatman GF/C glass fibre filters (5 cm diameter) that had previously been dried at 105°C for 2 h, cooled in a desiccator and weighed. Following filtration, the algae-loaded filters were again dried, cooled and weighed. The biomass of the suspension was calculated as mg DW l⁻¹. Owing to the high salt content in the NaCl-treated cultures, their DW was adjusted according to a factor determined by filtering a similar volume of 36 g l⁻¹ NaCl solution. These measurements were used to determine the brassinosteroid content in the samples.

Brassinosteroid quantification

The brassinosteroid content in the samples was analysed using the previously described method (Tarkowska *et al.*, 2016) with a few minor modifications. Briefly, 0.4–2.4 mg DW samples, each combined with 10 pmol of internal standard mixture comprising [²H₃]brassinolide, [²H₃]castasterone, [²H₃]24-*epi*brassinolide, [²H₃]24-*epi*castasterone, [²H₃]28-*nor*brassinolide, [²H₃]28-*nor*castasterone and [²H₃]typhasterol (OlChemim, Olomouc, Czech Republic), were extracted in ice-cold 60% acetonitrile overnight at 4°C with stirring. The crude extracts were subsequently centrifuged and the obtained supernatants were further purified on polyamine SPE columns (Supelco, Bellefonte, Pennsylvania, USA) and finally analysed by UHPLC-MS/MS (Waters, Milford, USA; Waters MS Technologies, Manchester, UK). Data analysis was carried out using Masslynx™ 4.1 software (Waters, Manchester, UK) and brassinosteroids were quantified by the standard isotope-dilution method using six technical replicates per one biological sample.




Fig. 1. Change in cell number in six microalgae exposed to high salt and cold temperature stress. Results are presented as mean \pm SE ($n = 6$). Different letters indicate significant differences ($P > 0.05$).

Results

Over the 7 h experimental period, three species i.e. *Chlorococcum ellipsoideum* (Fig. 1a), *Acutodesmus acuminatus* (Fig. 1d) and *Protococcus viridis* (Fig. 1e) continued to divide as indicated by a significant increase in cell number from 30 min to 7 h in the 25°C and 15°C treatments. Addition of NaCl had a negative effect on cell division with no significant changes in cell number over the 7 h period in the salt-stressed cultures. The other three species i.e. *Gyoeffya humicola* (Fig. 1b), *Nautococcus mamillatus* (Fig. 1c) and *Chlorella vulgaris* (Fig. 1f) did not undergo cell division over the 7 h experimental period with no significant change in cell number in the 25°C and 15°C treatments. Addition of NaCl stimulated cell division in *G. humicola* at 25°C (Fig. 1b) and *N. mamillatus* at 15°C (Fig. 1c).

Four endogenous brassinosteroids – BL, CS, homocasterone (homoCS) and typhasterol (TY) were detected in the six microalgae. In all species, regardless of the treatment, CS was present in the highest concentration. In *N. mamillatus*, *A. acuminatus*, *P. viridis* and *C. vulgaris* the other three

brassinosteroids were present in similar, lower amounts. In *G. humicola* homoCS occurred in moderate amounts with BL and TY present in the lowest concentrations, while in *C. ellipsoideum* BL and TY occurred in similar amounts with homoCS present in the lowest concentration (Supplementary table).

The total brassinosteroid content ranged from 52 to 670 pg mg⁻¹ DW, being lowest in *P. viridis* (Fig. 2e) and highest in *G. humicola* (Fig. 2b). Time of day and sample treatment had an effect on the brassinosteroid content. With the exception of *N. mamillatus* (Fig. 2c) and salt-stressed *C. ellipsoideum* at 25°C (Fig. 2a), the endogenous brassinosteroid content decreased over the course of the light period regardless of the treatment (Fig. 1).

The effect of salinity stress was pronounced, with generally higher brassinosteroid content in samples treated with 36 g l⁻¹ NaCl compared with the corresponding untreated cultures. The response to salinity stress was rapid, with samples harvested after 30 min exposure to high salinity generally having an increased brassinosteroid content compared with the corresponding untreated cultures (Fig. 2). In *G. humicola* and *A.*




Fig. 2. Endogenous brassinosteroid content in six microalgae exposed to high salt and low temperature stress. Results are presented as mean \pm SE (n = 6).

acuminatus, the content of brassinosteroids increased in the short term (30 min harvest) and then decreased over time (7 h harvest; Fig. 2b, d). Conversely, the brassinosteroid content increased more slowly in *C. ellipsoideum* in response to salinity stress when cultured at 25°C (Fig. 2a). In the other strains, the brassinosteroid content initially increased and remained at these elevated levels after 7 h.

In four strains, temperature had no effect on the brassinosteroid content while in *A. acuminatus* (Fig. 1d) and *C. vulgaris* (Fig. 1f), brassinosteroid levels decreased when transferred from 25 to 15°C. There was a salt-and-temperature accumulative effect in *G. humicola* where the brassinosteroid level was lower in the salt-stressed culture at 15°C (Fig. 2b) and higher in *N. mamillatus* (Fig. 2c).

Discussion

CS is the most widely distributed brassinosteroid in the plant kingdom, followed by BL with TY, 6-deoxocastasterone (6-deoxoCS), teasterone (TE) and 28-norcastasterone (28-norCS) also being commonly identified (Bajguz & Tretyn, 2003). The brassinosteroid profiles of the six microalgae analysed in the present study consisted of these common brassinosteroids with CS, BL, TY and homoCS being detected in all strains (Supplementary table). This is in agreement with previous studies where CS and BL were identified in 24 microalgae strains from diverse taxonomic divisions Chlorophyceae, Trebouxiophyceae, Ulvophyceae (all Chlorophyta) and one strain from the Streptophyta clade (Stirk *et al.*, 2013a); seven brassinosteroids, including CS, BL and TY as well as TE, 6-deoxoteasterone (6-deoxoTE), 6-deoxytyphasterol (6-deoxoTY) and 6-deoxoCS were isolated from *Chlorella vulgaris* (Bajguz, 2009a); two brassinosteroids with a C-24 configuration were present in the freshwater green alga *Hydrodictyon reticulatum* (Hydrodictyaceae; Yokota *et al.*, 1987); and CS and BL were isolated from the stipes and fronds of the kelp *Ecklonia maxima* (Stirk *et al.*, 2014).

In agreement with previous studies on non-vascular plants, the brassinosteroids identified in the six microalgae strains analysed in the present study are the last intermediates in the early C-6 oxidation pathway where TY is converted via CS to BL (Fujioka & Yokota, 2003). Feeding experiments where TE is converted via 3-dehydroteasterone to TY suggest that brassinosteroids are formed via the early C-6 oxidation pathway in the liverwort *Marchantia polymorpha* (Park *et al.*, 1999). This limited evidence suggests that the early C-6 biosynthetic pathway is the favoured pathway in non-vascular species although there is also evidence of the late C-6 oxidation pathway in *C. vulgaris* (Bajguz, 2009a). In contrast, the late C-6 oxidation pathway is the

predominate route in angiosperms such as *Arabidopsis*, tomato and tobacco (Fujioka & Yokota, 2003).

The brassinosteroid concentration ranges in the six strains analysed in the present study were similar to those recorded in the 24 microalgae strains previously analysed (118–978 pg mg⁻¹ DW; Stirk *et al.*, 2013a) but higher than reported in *C. vulgaris* (1.91 ng g⁻¹ DW; Bajguz, 2009a) and the kelp *Ecklonia maxima* (14–29 pg mg⁻¹ DW; Stirk *et al.*, 2014). The higher brassinosteroid content measured in the present study and the previously analysed 24 microalgae strains may be due to the cultures being harvested on day 4 while in an exponential growth phase. Brassinosteroids generally occur in higher concentrations in young, actively growing plant tissues compared with older tissues (Bajguz & Tretyn, 2003) as they are essential for cell elongation, influencing the mechanical properties of the cell wall during cell elongation (Bartwell *et al.*, 2013).

Of the four brassinosteroids detected in the six microalgae, CS, homoCS and TY have a 6-ketone B-ring and BL has a 7-oxolactone B-ring (Bajguz & Tretyn, 2003). In general, the 7-oxalactone brassinosteroids have the strongest biological activity followed by the 6-ketone brassinosteroids with the non-oxidized brassinosteroids having no activity (Bajguz & Tretyn, 2003). This has also been shown in microalgae where exogenous application of three BL and three CS analogues stimulated growth and increased protein and nucleic acid content in *C. vulgaris* with the three BL analogues being more effective than the CS analogues (BL > 24-*epi*BL > homoBL > CS > 24-*epi*CS > homoCS; Bajguz, 2000). In the present study, CS was consistently the predominant endogenous brassinosteroid in the six microalgae species, suggesting that brassinosteroids were actively involved in regulating physiological processes in the cells.

Previous studies investigating the role of brassinosteroids in growth and stress responses in microalgae used *C. vulgaris* as the model organism. Exogenous brassinosteroids were applied and changes in physiological processes and biochemical content measured. The present study is the first to report on changes in brassinosteroids in other microalgae species and to quantify changes in the endogenous brassinosteroids in response to external stimuli.

In the present study, addition of NaCl induced salt stress as seen by the decline in cell numbers in three species. There was a rapid response to salt stress with a significant increase in the endogenous brassinosteroid content within 30 min in all six species. The brassinosteroid content generally remained at elevated concentrations after 7 h in the light period. In contrast, a slower decrease in temperature from 25°C to 15°C had less effect on the brassinosteroid content

compared with the rapid salt-shock treatment in the six species. A similar rapid response was measured with exogenous application of brassinosteroids to heavy metal-stressed *C. vulgaris* where the synthesis of phytochelatins (organic ligands involved in chelation of heavy metals for detoxification) was rapid, with changes in the phytochelatin content occurring within the first few hours of brassinosteroid application (Bajguz, 2002).

In vascular plants, brassinosteroids are an integral part of the hormonal signalling network where they are actively involved in crosstalk with other phytohormones influencing certain physiological responses to environmental stimuli (Fariduddin *et al.*, 2014; Oklestková *et al.*, 2015; Rajewska *et al.*, 2016). In algae, crosstalk has been demonstrated in *C. vulgaris* where exogenous BL increased the thermotolerance of heat-stressed cultures by increasing the biosynthesis of abscisic acid (ABA) within 3 h of application (Bajguz, 2009b); exogenous BL also partially overcame the inhibitory effects of heavy metals on *C. vulgaris*, decreasing the accumulation of heavy metals in the cells and increasing ABA, indole-3-acetic acid (IAA) and zeatin content although there was no change in the endogenous BL content (Bajguz, 2011). Furthermore, brassinosteroids stimulated the levels of endogenous auxins in *C. vulgaris* and co-application of auxins and brassinosteroids acted synergistically to stimulate growth and the accumulation of proteins, chlorophylls and monosaccharides (Bajguz & Piotrowska-Niczyporuk, 2013); exogenous cytokinins stimulated the levels of endogenous brassinosteroids with a synergistic effect between exogenous cytokinins and exogenous BRs where there was an increase in cell number, proteins, chlorophylls and monosaccharides in a dose-effect relationship (Bajguz & Piotrowska-Niczyporuk, 2014).

The six microalgae strains analysed in the present study were part of a previous study where endogenous auxins, cytokinins, gibberellins and brassinosteroids were detected and quantified in 4-day old cultures (Stirk *et al.*, 2013a, 2013b). Thus it is likely that the rapid increase in endogenous brassinosteroids in response to salt stress in the present study would have been accompanied by changes in the other hormone concentrations. A more detailed investigation on the effects of external stimuli on hormone homeostasis is required in order to understand the role of hormones in regulating abiotic defence mechanisms in microalgae.

In conclusion, this was the first study to show that endogenous brassinosteroids increase in response to abiotic stress in a number of microalgae species. This response was rapid with increased brassinosteroid levels within 30 min of the salt shock. Understanding

the role of brassinosteroids in the stress responses of microalgae could enhance the value of *in vivo* production of bioactive microalgae metabolites.

Acknowledgements

This research was funded by the project SABANA (Grant no. 727874) from the European Union Horizon 2020 Research and Innovation Program, the National Research Foundation of South Africa (grant number 105997) and the Ministry of Education, Youth and Sports, Czech Republic (grant LO1204 from the National Program of Sustainability I and Agricultural Research).

Disclosure statement

No potential conflict of interest was reported by the authors.

Author contributions

WA Stirk: experimental design, wrote and edited manuscript; P Bálint: culture experiments; D Tarkowská: brassinosteroid analysis, drafting and editing manuscript; M Strnad: drafting and editing manuscript; J van Staden: drafting and editing manuscript; V Ördög: original concept; drafting and editing manuscript.

Supplementary information

The following supplementary material is accessible via the Supplementary Content tab on the article's online page at <https://doi.org/10.1080/09670262.2018.1441447>

Supplementary table. Brassinosteroids identified in the six microalgae strains exposed to salt and cold temperature stress and harvested after 30 min and 7 h exposure.

ORCID

Wendy A. Stirk  <http://orcid.org/0000-0003-4033-5641>

Danuše Tarkowská  <http://orcid.org/0000-0003-1478-1904>

Johannes van Staden  <http://orcid.org/0000-0003-0515-1281>

References

- Bajguz, A. (2000). Effect of brassinosteroids on nucleic acids and protein content in cultured cells of *Chlorella vulgaris*. *Plant Physiology and Biochemistry*, **38**: 209–215.
- Bajguz, A. (2002). Brassinosteroids and lead as stimulators of phytochelatins synthesis in *Chlorella vulgaris*. *Journal of Plant Physiology*, **159**: 321–324.
- Bajguz, A. (2009a). Isolation and characterization of brassinosteroids from algal cultures of *Chlorella vulgaris* Beijerinck (Trebouxiophyceae). *Journal of Plant Physiology*, **166**: 1946–1949.
- Bajguz, A. (2009b). Brassinosteroid enhanced the level of abscisic acid in *Chlorella vulgaris* subjected to short-term heat stress. *Journal of Plant Physiology*, **166**: 882–886.

- Bajguz, A. (2010). An enhancing effect of exogenous brassinolide on the growth and antioxidant activity in *Chlorella vulgaris* cultures under heavy metal stress. *Environmental and Experimental Botany*, **68**: 175–179.
- Bajguz, A. (2011). Suppression of *Chlorella vulgaris* growth by cadmium, lead and copper stress and its restoration by endogenous brassinolide. *Archives of Environmental Contamination and Toxicology*, **60**: 406–416.
- Bajguz, A. & Asami, T. (2004). Effects of brassinazole, an inhibitor of brassinosteroid biosynthesis, on light- and dark-grown *Chlorella vulgaris*. *Planta*, **218**: 869–877.
- Bajguz, A. & Czerpak, R. (1998). Physiological and biochemical role of brassinosteroids and their structure-activity relationship in the green alga *Chlorella vulgaris* Beijerinck (Chlorophyceae). *Journal of Plant Growth Regulation*, **71**: 131–139.
- Bajguz, A. & Piotrowska-Niczyporuk, A. (2013). Synergistic effect of auxin and brassinosteroids on the growth and regulation of metabolite content in the green alga *Chlorella vulgaris* (Trebouxiophyceae). *Plant Physiology and Biochemistry*, **71**: 290–297.
- Bajguz, A. & Piotrowska-Niczyporuk, A. (2014). Interactive effect of brassinosteroids and cytokinins on growth, chlorophyll, monosaccharide and protein content in the green alga *Chlorella vulgaris* (Trebouxiophyceae). *Plant Physiology and Biochemistry*, **80**: 176–183.
- Bajguz, A. & Tretyn, A. (2003). The chemical characteristic and distribution of brassinosteroids in plants. *Phytochemistry*, **62**: 1027–1046.
- Bajguz, A., Bajguz, A.J. & Tryniszewska, E.A. (2013). Recent advances in medicinal applications of brassinosteroids, a group of plant hormones. In *Studies in Natural Products Chemistry*, vol. 40 (Atta-ur-Rahman, F.R.S., editor), 33–49. Elsevier, Amsterdam.
- Bartwell, A., Mall, R., Lohani, P., Guru, S.K. & Arora, S. (2013). Role of secondary metabolites and brassinosteroids in plant defense against environmental stresses. *Journal of Plant Growth Regulation*, **32**: 216–232.
- Bold, H.C. (1949). The morphology of *Chlamydomonas chlamydogma* sp. nov. *Bulletin of the Torrey Botanical Club*, **76**: 101–108.
- Fariduddin, Q., Yusuf, M., Ahmad, I. & Ahmad, A. (2014). Brassinosteroids and their role in response of plants to abiotic stress. *Biologia Plantarum*, **58**: 9–17.
- Fujioka, S. & Yokota, T. (2003). Biosynthesis and metabolism of brassinosteroids. *Annual Review of Plant Biology*, **54**: 137–164.
- Oklestková, J., Rárová, L., Kvasnica, M. & Strnad, M. (2015). Brassinosteroids: synthesis and biological activities. *Phytochemical Reviews*, **14**: 1053–1072.
- Park, S.H., Han, K.S., Kim, T.W., Shim, J.K., Takatsuto, S., Yokota, T. & Kim, S.K. (1999). *In vivo* and *in vitro* conversion of teasterone to typhasterol in cultured cells of *Marchantia polymorpha*. *Plant and Cell Physiology*, **40**: 955–960.
- Rajewska, I., Talarek, M. & Bajguz, A. (2016). Brassinosteroids and response of plants to heavy metal action. *Frontiers in Plant Science*, **7**: 629.
- Stirk, W.A., Bálint, P., Tarkowská, D., Novák, O., Strnad, M., Ördög, V. & van Staden, J. (2013a). Hormone profiles in microalgae: Gibberellins and brassinosteroids. *Plant Physiology and Biochemistry*, **70**: 348–353.
- Stirk, W.A., Ördög, V., Novák, O., Rolčík, J., Strnad, M., Bálint, P. & van Staden, J. (2013b). Auxin and cytokinin relationships in 24 microalgal strains. *Journal of Phycology*, **49**: 459–467.
- Stirk, W.A., Tarkowská, D., Turečová, V., Strnad, M. & van Staden, J. (2014). Abscisic acid, gibberellins and brassinosteroids in Kelpak®, a commercial seaweed extract made from *Ecklonia maxima*. *Journal of Applied Phycology*, **26**: 561–567.
- Takatsuto, S., Abe, H. & Gamoh, K. (1990). Evidence for brassinosteroids in strobilus of *Equisetum arvense* L. *Agricultural and Biological Chemistry*, **54**: 1057–1059.
- Tarkowská, D., Novák, O., Oklestková, J. & Strnad, M. (2016). The determination of 22 natural brassinosteroids in a minute sample of plant tissue by UHPLC-MS/MS. *Analytical and Bioanalytical Chemistry*, **408**: 6799–6812.
- Yokota, T., Kim, S.K., Fukui, Y., Takahashi, N., Takeuchi, Y. & Takematsu, T. (1987). Brassinosteroids and sterols from a green alga, *Hydrodictyon reticulatum*: configuration at C-24. *Phytochemistry*, **26**: 503–506.